

Kynurenines IHC/IF Starter Pack

Ref: KY-ABSP

Test ImmuSmol's monoclonal and polyclonal antibodies against kynurenine pathway metabolites. For each antibody, IHC/IF validation pictures and technical specifications are provided: [visit individual product pages](#).

To make you own starter pack, please select 3 antibodies within our range :

INFORMATIONS

Product overview

Product name	Kynurenine pathway metabolites - antibody starter pack for IHC/IF
Immunogen	Conjugated amino acid metabolites
Clonality	Mouse monoclonal or rabbit polyclonal antibodies
Specificity	Each antibody was tested for affinity and specificity by competitive ELISA. ELISA data are available on product pages
Specie reactivity	Reacts with all species
Format	3 x 25µl

Reconstitution & storage

Form	Lyophilized powder
Storage	Store at 4°C
Storage buffer	Before use, vial should be resuspended in 25 µL of ultrapure water. Store at +4°C for short term (1-2 months). Aliquot and store at -20°C for long term. Avoid repeated freeze / thaw cycles

PROTOCOLS

Immunohistochemistry (IHC)	Dilute at 1:200-1:2000. Perform heat antigen retrieval before initiating IHC staining protocol on paraffin-embedded and frozen sections
Immunofluorescence (IF)	Dilute at 1:100-1:1000 on paraffin-embedded and frozen sections. Perform heat antigen retrieval and incubate with fluorescent dye-conjugated secondary antibody
Comments	Optimal working dilutions must be determined by the end-user
Restrictions	For research use only


REFERENCES

Antibodies are not yet cited. Submit an article and [get a 10% discount](#).


Selected publications on kynurenine pathway metabolites:

- [Hoshi M, Matsumoto K, Ito H, Ohtaki H, Arioka Y, Osawa Y, Yamamoto Y, Matsunami H, Hara A, Seishima M, Saito K. L-tryptophan-kynurenine pathway metabolites regulate type I IFNs of acute viral myocarditis in mice. J Immunol. 2012 Apr 15;188\(8\):3980-7. doi: 10.4049/jimmunol.1100997. Epub 2012 Mar 14.](#)
- [Berthon C, Fontenay M, Corm S, Briche I, Allorge D, Hennart B, Lhermitte M, Quesnel B. Metabolites of tryptophan catabolism are elevated in sera of patients with myelodysplastic syndromes and inhibit hematopoietic progenitor amplification. Leuk Res. 2013 May;37\(5\):573-9. doi: 10.1016/j.leukres.2013.02.001. Epub 2013 Feb 28.](#)
- [Schwarcz R, Bruno JP, Muchowski PJ, Wu HQ. Kynurenines in the mammalian brain: when physiology meets pathology. Nat Rev Neurosci. 2012 Jul;13\(7\):465-77. doi: 10.1038/nrn3257.](#)
- [Pilotte L, Larrieu P, Stroobant V, Colau D, Dolusic E, Frédérick R, De Plaen E, Uyttenhove C, Wouters J, Masereel B, Van den Eynde BJ. Reversal of tumoral immune resistance by inhibition of tryptophan 2,3-dioxygenase. Proc Natl Acad Sci U S A. 2012 Feb 14;109\(7\):2497-502. doi: 10.1073/pnas.1113873109. Epub 2012 Jan 30.](#)

Product pictures




12 validated antibodies, from L-Tryptophan to downstream kynurenine metabolites




L-Kynurenine detection by IHC in human colon tumor (mouse mAb)



Limited stock offer: Try 3 antibodies for the price of 1



Kynurenic acid detection in human brain by IF (mouse mAb)



Quinolinic acid detection in human parkinson's brain (rabbit pAb)

Contact information

Immusmol
229 Cours de l'Argonne
33 000 Bordeaux - France
Tel: +33 (0) 5 6431 1170
www.immusmol.com

To order, review, ask for technical support, visit product page at:

<https://www.immusmol.com/shop/kynurenines-ihc-if-starter-pack/>